

**δ –invariant for Quasi-periodic Oscillations and Physical
Parameters of 4U 0614+09 binary**

Lev Titarchuk

NASA/ Goddard Space Flight Center, Greenbelt MD 20771, and George Mason
University/CSI, USA; lev@lheapop.gsfc.nasa.gov

Vladimir Osherovich

NASA/Goddard Space Flight Center/RITSS, Greenbelt MD 20771 USA;
vladimir@urap.gsfc.nasa.gov

Received _____; accepted _____

ABSTRACT

The recently formulated Two Oscillator (TO) model interprets the lowest of the kilohertz frequencies of the twin peak quasi-periodic oscillations in X-ray binaries as the Keplerian frequency ν_K . The high twin frequency ν_h in this model holds the upper hybrid frequency relation to the rotational frequency of the neutron star’s magnetosphere Ω : $\nu_h^2 = \nu_K^2 + 4(\Omega/2\pi)^2$. The vector Ω is assumed to have an angle δ with the normal to the disk. The first oscillator in the TO model allows one to interpret the horizontal branch observed below 100 Hz as the lower mode of the Keplerian oscillator under the influence of the Coriolis force, with frequency ν_L being dependent on ν_h , ν_K and δ . For some stars such as 4U 0614+09, Sco X-1 and 4U 1702-42, ν_h , ν_K and ν_L have been observed simultaneously providing the opportunity to check the central prediction of the TO model: the constancy of δ for a particular source. Given the considerable variation of each of these three frequencies, the existence of an observational invariant with a clear physical interpretation as a global parameter of the neutron star magnetosphere is an important test of the TO model. Using the results of recent observations of 4U 0614+09 we verify the existence of this invariant and determine the angle $\delta = 15^\circ.6 \pm 0.^{\circ}5$ for this star. The second oscillator in the model deals with a radial (presumably sound) oscillation and diffuse process in the viscous layer surrounding the neutron star. Our analysis of the viscous oscillation frequency ν_V and the break frequency ν_b of the diffusion shows that the spin value of the inner boundary of the transition layer for 4U 0614+09 is at least two times more than values for 4U 1728-34 and Sco X-1.

Subject headings: accretion, accretion disks—diffusion—stars:individual (4U 0614+09, 4U 1728-34, Sco X-1, 4U 1702-42)—stars:neutron—X-ray:star—waves

1. Introduction

This *Letter* contains further verification of the Two Oscillator (TO) model for kHz QPO's in NS binaries and the related low frequency features using the observational results for 4U 0614+09 from van Straaten et al. (2000) (hereafter VS00). In a series of the papers (Osherovich & Titarchuk 1999a, Titarchuk & Osherovich 1999, Osherovich & Titarchuk 1999b, Titarchuk, Osherovich & Kuznetsov 1999, hereafter OT99a, TO99, OT99b and TOK respectively) the authors offered the TO model which explained the variation of the frequency difference between two kHz QPOs suggesting that the upper one, namely ν_h is an upper hybrid frequency of the Keplerian oscillator under the influence of the Coriolis force and *the lower kHz QPO is the Keplerian frequency ν_K* . This identification is a principal difference between the TO model and other QPO models presented so far (e.g. Stella & Vietri 1998 Miller, Lamb, Cook 1998, Kaaret, Ford & Chen 1997) since in other models, the highest kHz QPO is identified as a Keplerian frequency.

According to the TO model, the Keplerian oscillator has two branches characterized by high frequency ν_h (~ 1 kHz) and by low frequency ν_L ($\sim 50 - 100$ Hz). The frequency ν_L depends strongly on the angle δ between the normal to the neutron star disk and $\mathbf{\Omega}$ —the angular velocity of the magnetosphere surrounding the neutron star. In the lower part of the QPO spectrum (~ 10 Hz), the second oscillator of the TO model describes the physics of the viscous transition layer, namely, radial viscous oscillations with frequency ν_V (previously called “extra noise component”) and the diffusive process in the transition region (the innermost part of the disk) which is characterized by the break frequency ν_b . According to the TO model, all frequencies (namely ν_h , ν_L , ν_b and ν_V) have specific dependences on ν_K . Correlations of ν_b and ν_V with kHz frequencies are fitted in TO99 by the theoretical curves using the dimensionless parameter, a_K :

$$a_K = m(x_0/3)^{3/2}(\nu_0/363 \text{ Hz}) \quad (1)$$

where normalized mass $m = M/M_\odot$, radius of the inner edge of disk $x_0 = R_0/R_S$ in Schwarzschild radius (R_S units) and ν_0 is the spin of the disk inner edge.

Van Straaten et al. (2000) have found for 4U 0614+09 that dependences of ν_V and ν_b on ν_K do not follow the theoretical curves which fit the data points for 4U1728-34 (TO99) and Sco X-1 (TOK). In this *Letter*, we revisit this issue and show that the observational correlations for source 4U 0614+09 are related to the theoretical ones for the same parameter $a_K = 1.03$ (as for 4U 1728-34 and Sco X-1) but with a spin of the inner edge of accretion disk higher than 363 Hz. Such a high spin leads to the high compactness of the central object because

$$R_0 = 9 \text{ km} \times m^{1/3} \left(\frac{a_K}{\nu_0/363 \text{ Hz}} \right)^{2/3}. \quad (2)$$

We check predictions of the TO model regarding ν_V vs ν_K dependence as well as relation between ν_b and ν_K for all available observations (see §3).

In OT99b, the authors demonstrated for the source 4U 1702-42 that the inferred angle δ , (see Eq 2, 5 there)

$$\delta = \arcsin \left[(\nu_h^2 - \nu_K^2)^{-1/2} (\nu_L \nu_h / \nu_K) \right] \quad (3)$$

depends only, on ν_K , ν_h and ν_L and stays the same ($3.9^\circ \pm 0.2^\circ$) over significant range of ν_K (650-900 Hz). In Formula (3), δ is assumed to be small. A general formulation for any δ is presented, in OT99a with conjecture that eventually δ as an invariant will be found for all 20 sources known to have QPOs. Using series of observations where ν_h , ν_K , ν_L are detected simultaneously (van der Klis et al. 1997; Markwardt, Strohmayer & Swank 1999; van Straaten et al. 2000) we check that the angle δ is a true invariant for three specific sources: Sco X-1, 4U 1702-42, 4U 0614+09 (see §2). We will also present the magnetospheric rotational profile for 4U 0614+09 inferred from VS00 data in §4. Discussion and summary follow in the last section.

2. δ –Invariant and Verification of Two-Oscillator Model

We have used the frequencies measured for sources 4U 0614+09, Sco X-1 and 4U 1702-42 where the low and high kHz QPO peaks ν_K and ν_h and HBO frequencies ν_L are measured simultaneously. The resulting values of δ calculated from Eq. (3) are shown in Figure 1. Indeed, for each of these sources, the δ -values show little variation with ν_K , ν_h , ν_L . They are $3^\circ.9 \pm 0.2$ and $5^\circ.5 \pm 0.5$ for 4U 1702-42 and Sco X-1 respectively. For the source 4U 0614+09 the point at $\nu_K = 821.6$ Hz is not included since the ν_K peak is very broad with FWHM= 205 Hz, which is almost order of magnitude larger than that for any other points for the same star according to Table 1 in VS00. The angle δ obtained for the source 4U 0614+09 is $15^\circ.6 \pm 0^\circ.5$ and is significantly larger than for other stars. The existence of the δ -invariant predicted by the TO model (OT99b) is a challenge for any other QPO model. It is important to note that *all frequencies included in this δ -relation are observed frequencies and thus the relation Eq. (3) is a model independent invariant.*

3. Break Frequency vs kHz QPO Frequency Correlations

Further tests of the TO model can be done using a comparison of the observed correlation of break and kHz QPO frequencies (VS00) with the theoretical dependences derived in Titarchuk, Lapidus & Muslimov (1998, hereafter TLM) and TO99. In Fig. (2) (upper panel), we present the theoretical curves calculated using Eq. (9) in TO99 for different values of $\nu_{0,363} = \nu_0/363$ Hz. For the source 4U 0614+09 we found the Lebesgue’s measure (for definition, see TOK) is better than 15% for $a_K = 1.03$ and $\nu_{0,363} > 2$ which is at least 3 times less than that for $\nu_{0,363} = 1$ (see upper panel in Fig. 2). We fitted the data points by the curves with $\nu_{0,363} > 3$ ($\nu_0 > 1089$ Hz) and we have found that the quality of those fits is improved only slightly. Thus it is difficult to get the constraints on ν_0 better than $\nu_0 \gtrsim 800$ Hz. Recently Bhattacharyya et al. (2000) using the full General Relativity technique argued that the NS in Cyg X-2 rapidly rotates with the frequencies of order of 1 kHz which is very close to that we obtained here for 4U 0614+09.

We approximate the solution of Eq (9) of TO99 by a polynomial for $a_K = 1.03$ and $\nu_{0,363} = 1$, namely,

$$\nu_b = C_b \cdot P_4(\nu_K) = C_b(B_1\nu_K + B_2\nu_K^2 + B_3\nu_K^3 + B_4\nu_K^4) \quad (4)$$

where $B_1 = -1.68 \times 10^{-4}$, $B_2 = 1.36 \times 10^{-5} \text{ Hz}^{-1}$, $B_3 = -2.47 \times 10^{-8} \text{ Hz}^{-2}$, $B_4 = 4.45 \times 10^{-11} \text{ Hz}^{-3}$. We fit the observational break frequencies by the theoretical curve which is a function of parameter $\nu_{0,363}$. For a given $\nu_{0,363}$ that curve can be obtained by the scaling the argument x of the polynomial $P_4(x)$:

$$\nu_b(\nu_K) = C_b P_4(\nu_K/\nu_{0,363}). \quad (5)$$

The normalization of the curve is controlled by the constant C_b which reflects the properties of the specific source. For 4U 1728-34 the observed frequencies are fitted by the curve with $a_K = 1.03$ (TO99) for which the constant $C_b = 1$ with $R_0 = 11 \text{ km}$ for 1.4 solar masses. For 4U 0614+09 we fit the data points by the curves with the parameters $\nu_{0,363} = 1, 2, 3$, (i.e. $\nu_0 = 363 \text{ Hz}$, 726 Hz and 1089 Hz) for which the corresponding constants $C_b = 2, 22.8, 55.4$ are chosen respectively. Using Eq (2), one can estimate mass-radius (MR) relations for a given value of $\nu_{0,363}$. For $\nu_{0,363} = 2$ and 3 the radius $R_0 \approx 6.4$ and 4.9 km respectively, for the standard NS mass, 1.4 solar masses. There is definitely an indication of higher compactness for 4U 0614+09. It is true that there is a possibility to elaborate a much more stringent condition on the MR relation using Eq. (2) and fits of the data in Fig. 2 which may lead to small masses and radii for the star – $m < 1$ and $R < 9 \text{ km}$. In that case such a low mass compact star can be easily explained in terms of strange star models (e.g. Bombaci 1997). In fact, under constraints imposed by the TO model Li et al. 1999 pointed out the possibility that 4U 1728-34 is a strange star rather than a neutron star. Thus our arguments of the high compactness in 4U 0614+09 support a strange star concept. But the final conclusion on MR constraints cannot be worked out without the full relativistic treatment which is out of the scope of this paper.

According to the low panel in Fig. 2, ν_h is not lower than 400 Hz for any value of ν_K . Because

$$\nu_h = [\nu_K^2 + 4(\Omega/2\pi)^2]^{1/2}, \quad (6)$$

the rotational magnetospheric frequencies $\nu_{mag} = (\Omega/2\pi)$ should have an asymptote for large radii (or small values $\nu_K \lesssim 2\nu_{mag}$) which equals approximately 200 Hz. The detailed profile of ν_{mag} constructed below confirms this expectation.

4. Inferred Rotational Frequency Profile of the NS Magnetosphere

From the observed kHz frequencies (VS00, Table 1), ν_h and ν_K , the profile of $\nu_{mag} = \Omega(\nu_K)/2\pi$ has been calculated according to formula (6) and modeled using theoretically inferred magnetic multipole structure of differentially rotating magnetosphere (OT99a)

$$\nu_{mag} = C_0 + C_1\nu_K^{4/3} + C_2\nu_K^{8/3} + C_3\nu_K^4 \quad (7)$$

where $C_2 = 2(C_1C_3)^{1/2}$. The constants $C_0 \equiv \nu_{mag}^0 = 190$ Hz, $C_1 = 6.52 \times 10^{-2}$ Hz $^{-1/3}$, $C_2 = -1.19 \times 10^{-5}$ Hz $^{-5/3}$ and $C_3 = 0.91 \times 10^{-9}$ Hz $^{-3}$ have been obtained by the least-squares fit with $\chi^2 = 29.8/25$ (OT99a). The resulting fit is shown in the lower panel of Fig. 3 by solid line. The ν_{mag} profiles for Sco X-1 (which is very similar to those for 4U 1608-52 and 4U 1728-34) and that for 4U 0614+09 differ in the sign of curvature: it is negative for the profile of Sco X-1 profile and positive for that of 4U 0614+09. The ν_{mag} profiles for Sco X-1, 4U 1608-52 (as well as for 4U 1728-34) according to analytical formula (14) in OT99a have a maximum at large ν_K (small radii) and also have a minimum at small ν_K (large radii). The adjustment of the Keplerian disk to the sub-Keplerian rotation of the NS in 4U 0614+09 occurs at much larger distances from the central source (3.6-7 NS radii) than that for 4U 1728-34 where the adjustment radii are within 1.6-3.4 NS radii. Thus the blobs (formed at the adjustment radius as result of the super-Keplerian motion [TLM]) presumably probe the different parts of the magnetosphere: in the 4U 0614+09 case, it is

the outer part of the magnetosphere, near the minimum of the rotational frequency while in the 4U 1728-34 case, it is a part which is closer to the inner part, near the maximum. Thus one should make an expansion of ν_{mag} over the Chandrasekhar potential A near the minimum in the former case and near the maximum in the latter case (see OT99a for details). We may try to predict ν_{mag} for much smaller ν_K (around 150 Hz) than those frequencies ν_K (between 400 and 850 Hz) where we have constructed the profile (7). Using VS00 data points for 8 August with $\nu_K = 153.4 \pm 5.6$ Hz and $\nu_h = 449.4 \pm 19.5$ Hz and Eq. (6), we find $\nu_{mag} = 211. \pm 23.$ Hz which is in good agreement with the value $\nu_{mag} = 235$ Hz obtained from the analytical expression, Eq. (7).

As shown in Fig. 3 (the upper panel) the difference in ν_{mag} profiles for Sco X-1 and for 4U 0614+09 results in different profiles for $\Delta\nu = \nu_h - \nu_K$: namely, $\Delta\nu$ monotonically decreases with ν_K for Sco X-1 but has a minimum for 4U 0614+09. According to VS00 $\Delta\nu = 311.8 \pm 1.8$ Hz except one point (not included in Fig. 3) which deviates 11σ from this trend. We will return to this point in another paper.

5. Break Frequency vs Low Lorentzian QPO Frequency Correlation

According to solutions presented in TO99 and TOK, the break frequency ν_b correlates with the low Lorentzian frequency ν_V (which we interpreted as a viscous frequency in the TO model) through power law relation

$$\nu_b \approx 0.041\nu_V^{1.61}. \quad (8)$$

It was confirmed in TOK using the data of Wijnands & van der Klis (1999) that the same index approximately is valid for the wide class of NS and BH objects. In Figure 4 we demonstrate that the observed correlation of ν_b vs ν_V in 4U 0614+09 is described well by the same power law dependence and the quality of this fit is similar to that of 4U 1728-34, i.e. the Lebesgue's measure is less than 20%. Equations (5) and (8) give us theoretical

dependences of ν_V on ν_K , assuming $\nu_{0,363} = 3$. For August 8, 1996 the theoretical value $\nu_V \approx 9$ Hz with inherited 20% accuracy comes close to the observed $\nu_V = 11.2 \pm 1.4$ Hz as given in VS00.

6. Discussion and Conclusions

In this *Letter* we have verified predictions of the two-oscillator (TO) model for three sources. The unifying characteristic of spectra for both oscillators which share the common boundary at the outer edge of the viscous transition layer is the strong dependence of all frequencies in the model on ν_K . The angle δ as a global parameter describes the inclination of the magnetospheric equator to the equatorial plane of the disk. Measured locally for different radial distances (therefore different ν_K), δ may vary considerably if the observed oscillations do not correspond to the predicted low Keplerian branch. The constancy of δ shown in Figure 1 argues in favor of the TO model. The parameter δ is critical in evaluating the differences between the spectra of different sources and we suggest that the higher ν_L for 4U 0614+09 and 4U 1728-34 (than that for Sco X-1) are mainly a result of a larger angles δ .

We consider the angle δ to be a fundamental geometric parameter for neutron stars and may be related to the evolution and prehistory of these systems. During the NS life time, the NS is believed to spin up gaining more angular momentum from the companion and the axis of magnetosphere may change with respect the rotational axis of the star (or the disk) (Bhattacharya 1995). The X-ray bursters are found in old population (age of order 10^9 years) concentrated towards the galactic center (Lewin, van Paradijs & Taam 1995) and thus it is not surprising that in these systems the angle δ between spin axis and the magnetic axis is small presumably as a result of long evolution. The absence of X-ray pulsations from the LMXBs and the fact that type I X-ray bursters never occur in systems which show pulsations (Lewin, van Paradijs & Taam 1995) probably can be explained by

the small angles δ . In contrast, in X-ray pulsars which are mostly observed in the High Mass Binaries (HMXBs), δ is not small: the mean value of δ is 27° (Leahy 1991) and the age of these systems is much smaller, of order of 10^7 years.

The authors acknowledge discussions with Joe Fainberg, Sergey Kuznetsov and Craig Markwardt. We are grateful to Steve van Straaten for the data which enabled us to make comparisons with the data. We are also grateful the referee for the fruitful suggestions.

REFERENCES

- Bhattacharya, D. (1995), X-ray Binaries (Eds. Lewin, W. H.G. van Paradijs, J. & van den Heuvel, E.P.J.) (Cambridge: University Press) p.233
- Bhattacharyya, S., Thampan, A.V., Misra, R., & Datta, B. 2000, ApJ, in press
- Bombaci, I. 1997, Phys. Rev. C 55, 1587
- Kaaret, P., Ford, E.C., & Chen, K. 1997, ApJ, 480, L27
- Li, X.D., Ray, S., Dey, J., Dey, M., & Bombaci, I. 1999, ApJ, 527, L51
- Lewin, W. H.G. van Paradijs, J. & Taam, R. (1995), X-ray Binaries (Eds. Lewin, W. H.G. van Paradijs, J. & van den Heuvel, E.P.J.) (Cambridge: University Press) p.175
- Leahy, D.A. 1991, MNRAS, 251, 203
- Markwardt, C.B., Strohmayer, T.E., & Swank, J.H. 1999 ApJ, 512, L125
- Miller, M.C., Lamb, F.K., & Cook, D. 1998, ApJ, 508, 791
- Osherovich, V.A. & Titarchuk, L.G. 1999, ApJ, 522, L113 (OT99a)
- Osherovich, V.A. & Titarchuk, L.G. 1999, ApJ, 523, L73 (OT99b)
- Stella, L., & Vietri, M. 1998, ApJ, 492, L59
- Strohmayer, T. E., Zhang, W., Swank, J. H., Smale, A., Titarchuk, L., Day, C., & Lee, U. 1996, ApJ, 469, L9
- Titarchuk, L., & Osherovich, V., & Kuznetsov, S. 1999, ApJ, 525, L129 (TOK)
- Titarchuk, L., & Osherovich, V. 1999, ApJ, 518, L95 (TO99)
- Titarchuk, L., Lapidus, I.I., & Muslimov, A. 1998, ApJ, 499, 315 (TLM)

van der Klis, M., Wijnands, R.A.D., Horne, K. & Chen, W. 1997, ApJ, 481, L97

van der Klis, M. *et al.* 1996, ApJ, 469, L1

van Straaten, S., Ford, E., van der Klis, M., Mendez, M., & Kaaret, Ph. 2000, ApJ, in
press, astro-ph/0001480 (VS00)

Wijnands, R. A.D. & van der Klis, M., 1999, ApJ, 514, 939

Fig. 1.— δ -invariant. Inferred angles δ between the rotational axis of magnetosphere and the normal to the plane of Keplerian oscillations as a function of low peak kHz QPO frequency for three sources: 4U 0614+09, Sco X-1, and 4U 1702-42. The δ -invariant (eq.[3]) is calculated using the observational frequencies ν_h , ν_K and ν_L only.

Fig. 2.— *Low panel:* Break frequency of a broken power law function of the power spectrum density versus the frequency of the kHz QPOs for 4U 0614+09 (Straaten et al. 2000). Open circles are for the ν_b -dependence versus low kHz peak and open squares are the ν_b -dependences versus high kHz peak. The solid line is the theoretical curve for break frequencies calculated using (TO99 eqs. [4-5, 9, 13] corresponding to the parameter $a_k = 1.03$ and $\nu_0 = 1089$ Hz. *Upper panel:* The theoretical dependences of the break frequency versus the low peak kHz QPOs for different values of the spin frequency of the inner edge of accretion disk ν_0 . Observational points for 4U 0614+09 (Straaten et al. 2000) *open circles* and for 4U 1728-34 (Ford & van der Klis 1998) *solid circles*

Fig. 3.— *Low panel:* Inferred rotational frequency of the NS magnetosphere (solid line) as a function of the frequency of the low kHz QPO peak (Eq. 7 for 4U 0614+09 and see OT99a for Sco X-1). For 4U 0614+09 (van Straaten et al. 2000) observational points shown as *solid circles* and for Sco X-1 (van der Klis et al. 1997) as *open circles*. *Upper panel:* The peak difference of kHz QPO frequencies for the same sources. The solid lines are theoretical curves obtained using the above theoretical dependences of the rotational frequencies.

Fig. 4.— Break frequency versus main viscous frequency for 4U 0614+09. Solid line is the theoretical curve which is almost a power law with index 1.61.







